

Baryonic Isgur-Wise functions in large- N_c heavy quark effective theory

Ming-Kai Du* and Chun Liu†

 State Key Laboratory of Theoretical Physics, Institute of Theoretical Physics, Chinese Academy of Sciences,
 P.O. Box 2735, Beijing 100190, China

(Received 23 March 2013; published 15 May 2013)

Large- N_c relations among baryonic Isgur-Wise functions appearing at the order of $1/m_Q$ are analyzed. An application to $\Omega_b \rightarrow \Omega_c$ weak decays is given.

DOI: 10.1103/PhysRevD.87.094015

PACS numbers: 12.39.Hg, 12.38.Lg

I. INTRODUCTION

Weak decays of heavy baryons are interesting both experimentally and theoretically. They are now under the study of the LHC experiments, as well as previous Tevatron and LEP experiments. They also provide a playing ground for nonperturbative QCD methods. Heavy baryons containing a single heavy quark are described by the heavy quark effective theory (HQET) [1–4]. Relevant physical quantities can be factorized into a calculable perturbative part and universal hadronic quantities. To calculate the latter, some nonperturbative methods, like the large- N_c one [5], are needed.

Consider the heavy baryon weak transitions $\Lambda_b \rightarrow \Lambda_c$ and $\Sigma_b^{(*)} \rightarrow \Sigma_c^{(*)}$. The matrix elements of vector and axial currents ($V^\mu = \bar{c}\gamma^\mu b$ and $A^\mu = \bar{c}\gamma^\mu\gamma^5 b$) between the Λ_b and Λ_c can be parametrized as

$$\begin{aligned} \langle \Lambda_c(v', s') | V^\mu | \Lambda_b(v, s) \rangle &= \bar{u}_{\Lambda_c}(v', s') (F_1(\omega)\gamma^\mu + F_2(\omega)v^\mu \\ &\quad + F_3(\omega)v'^\mu) u_{\Lambda_b}(v, s), \\ \langle \Lambda_c(v', s') | A^\mu | \Lambda_b(v, s) \rangle &= \bar{u}_{\Lambda_c}(v', s') (G_1(\omega)\gamma^\mu + G_2(\omega)v^\mu \\ &\quad + G_3(\omega)v'^\mu)\gamma^5 u_{\Lambda_b}(v, s), \end{aligned} \quad (1)$$

and those between Σ_b and $\Sigma_c^{(*)}$ are

$$\begin{aligned} \langle \Sigma_c(v', s') | V^\mu | \Sigma_b(v, s) \rangle &= \bar{u}_{\Sigma_c}(v', s') (F'_1\gamma^\mu + F'_2v^\mu + F'_3v'^\mu) u_{\Sigma_b}(v, s), \\ \langle \Sigma_c(v', s') | A^\mu | \Sigma_b(v, s) \rangle &= \bar{u}_{\Sigma_c}(v', s') (G'_1\gamma^\mu + G'_2v^\mu + G'_3v'^\mu)\gamma^5 u_{\Sigma_b}(v, s), \\ \langle \Sigma_c^*(v', s') | V^\mu | \Sigma_b(v, s) \rangle &= \bar{u}_{\Sigma_c^*}(v', s') (N'_1v^\lambda\gamma^\mu + N'_2v^\lambda v^\mu + N'_3v^\lambda v'^\mu \\ &\quad + N'_4g^{\lambda\mu})\gamma^5 u_{\Sigma_b}(v, s), \\ \langle \Sigma_c^*(v', s') | A^\mu | \Sigma_b(v, s) \rangle &= \bar{u}_{\Sigma_c^*}(v', s') (K'_1v^\lambda\gamma^\mu + K'_2v^\lambda v^\mu + K'_3v^\lambda v'^\mu \\ &\quad + K'_4g^{\lambda\mu}) u_{\Sigma_b}(v, s), \end{aligned} \quad (2)$$

where $\omega = v \cdot v'$ and $F_i(\omega)^{(l)}$, $G_i(\omega)^{(l)}$, $N_i(\omega)'$, and $K_i(\omega)'$ are general form factors. As is well known, in the HQET, form factors can be described in terms of several independent universal form factors which are the so-called Isgur-Wise functions.

The Isgur-Wise functions are defined as follows. Note that it is self-evident that in the HQET, heavy quark fields and baryon fields have their own definition, in spite of adopting the same symbols as in full QCD. For the $\Lambda_b \rightarrow \Lambda_c$ transition, at the leading order of heavy quark expansion, there is only one Isgur-Wise function $\eta(\omega)$,

$$\langle \Lambda_c(v', s') | \bar{c}\Gamma b | \Lambda_b(v, s) \rangle = \eta(\omega) \bar{u}_{\Lambda_c}(v', s') \Gamma u_{\Lambda_b}(v, s), \quad (3)$$

where Γ stands for general γ matrices, and $\eta(\omega)$ is normalized at the zero recoil, namely, $\eta(1) = 1$. For $\Sigma_b^{(*)} \rightarrow \Sigma_c^{(*)}$ transitions, two Isgur-Wise functions $\xi_1(\omega)$ and $\xi_2(\omega)$ appear at the leading order [6],

$$\begin{aligned} \langle \Sigma_c^{(*)}(v', s') | \bar{c}\Gamma b | \Sigma_b^{(*)}(v, s) \rangle \\ = [-g_{\mu\nu}\xi_1(\omega) + v_\mu v'_\nu \xi_2(\omega)] \bar{u}_{\Sigma_c^{(*)}}^\mu(v', s') \Gamma u_{\Sigma_b^{(*)}}^\nu(v, s), \end{aligned} \quad (4)$$

where $\xi_1(1) = 1$, and $u_{\Sigma_c^{(*)}}^\nu$ is the Rarita-Schwinger spinor for a spin- $\frac{3}{2}$ particle. And $u_{\Sigma_b^{(*)}}^\nu$ is defined as

$$u_{\Sigma_b^{(*)}}^\nu(v, s) = \frac{\gamma^\nu + v^\nu}{\sqrt{3}} \gamma_5 u_{\Sigma_b}(v, s). \quad (5)$$

Isgur-Wise functions at $1/m_Q$ order will be discussed in detail in the next section.

Then, in the heavy quark limit, the general form factors in Eqs. (1) and (2) are simplified. For the $\Lambda_b \rightarrow \Lambda_c$ transition,

$$F_1 = G_1 = C(\mu)\eta(\omega), \quad F_2 = G_2 = F_3 = G_3 = 0, \quad (6)$$

where $C(\mu)$ is a perturbatively calculable coefficient. For $\Sigma_b^{(*)} \rightarrow \Sigma_c^{(*)}$ transitions, the formulas are a bit more complex, which can be found in [7]. Note that Isgur-Wise functions are independent of the weak currents.

At this stage, the Isgur-Wise functions are still unknown and need nonperturbative methods to be calculated. In the

*mkdu@itp.ac.cn
 †liuc@mail.itp.ac.cn

large- N_c limit, interesting information about baryonic Isgur-Wise functions was obtained. In large- N_c baryons, there is a spin-flavor symmetry of light quarks [8] which not only gives the mass degeneracy of $\Sigma_Q^{(*)}$ and Λ_Q , but also results in the following relations among the Isgur-Wise functions [9,10]:

$$\xi_1(\omega) = \eta(\omega), \quad \xi_2(\omega) = \frac{\eta(\omega)}{1 + \omega}. \quad (7)$$

This large- N_c result is also consistent with that obtained from the large- N_c constituent quark model [11].

Furthermore, in the heavy baryon Skyrme model [12], $\eta(\omega)$ is calculated to be [13]

$$\eta(\omega) = 0.99 \exp[-1.3(\omega - 1)]. \quad (8)$$

In fact, in the real large- N_c limit, $\eta(\omega)$ is actually a δ function [14].

From Eqs. (3), (4), and (7), it is observed that to obtain the $\Sigma_Q^{(*)}$ matrix elements of weak currents in the large- N_c approximation, what we need to do is just multiply the Λ_Q matrix element by the following Lorentz tensor:

$$\left[-g_{\mu\nu} + \frac{v_\mu v'_\nu}{1 + \omega} \right]. \quad (9)$$

This is because the two kinds of decays are essentially the same, except for Lorentz structures (a kinetic result of the light degrees of freedom).

With this observation, in the following sections, we will extend the large- N_c relations in Eq. (7) to $O(1/m_Q)$.

II. ISGUR-WISE FUNCTIONS AT $O(1/m_Q)$

There are more Isgur-Wise functions at the $1/m_Q$ order. Before we consider large- N_c relations among the Isgur-Wise functions at $O(1/m_Q)$, it is useful to start from their definition.

A. A mini-review

$1/m_Q$ corrections arise from two sources. One is due to the HQET Lagrangian at $O(1/m_Q)$, and the other is obtained through the $1/m_Q$ expansion of heavy quark currents in the full QCD.

Firstly, let us consider the Lagrangian corrections. To the order $O(1/m_Q)$ the effective Lagrangian is [2,15]

$$\mathcal{L} = \bar{h}_v i v \cdot D h_v - \frac{1}{2m_Q} \bar{h}_v \left[D^2 + \frac{1}{2} g_s \sigma_{\mu\nu} G^{\mu\nu} \right] h_v.$$

For the hadronic matrix element of a heavy quark current, correction due to the heavy quark kinetic energy is

$$\langle H_c | -i \int d^4 x T \left(g_s \bar{c}_{v'} \frac{D^2}{2m_c} c_{v'} |_x \bar{c}_{v'} \Gamma b_v |_0 \right) | H_b \rangle. \quad (10)$$

For H_Q being Λ_Q , Eq. (10) is parametrized as

$$\bar{u}_{\Lambda_c}(v', s') \frac{1 + \not{v}'}{2} \Gamma u_{\Lambda_b}(v, s) \frac{\chi(\omega)}{m_c}, \quad (11)$$

where $\chi(\omega)$ is the Λ_Q Isgur-Wise function at the order of $1/m_Q$. It satisfies that $\chi(1) = 0$. In the real large- N_c limit $\chi(\omega) = 0$ [14].

In the case of H_Q being $\Sigma_Q^{(*)}$, Eq. (10) is parametrized via two more Isgur-Wise functions,

$$\frac{1}{m_c} [-g_{\mu\nu} \chi_1(\omega) + v_\mu v'_\nu \chi_2(\omega)] \bar{u}_{\Sigma_c^{(*)}}^\mu(v', s') \times \frac{1 + \not{v}'}{2} \Gamma u_{\Sigma_b^{(*)}}^\nu(v, s), \quad (12)$$

with that $\chi_1(1) = \chi_2(1) = 0$.

Correction due to the heavy quark chromomagnetic interaction is

$$\langle H_Q | -\frac{i}{2} \int d^4 x T \left(g_s \bar{c}_{v'} \frac{\sigma_{\mu\nu} G^{\mu\nu}}{2m_c} c_{v'} |_x \bar{c}_{v'} \Gamma b_v |_0 \right) | H_Q \rangle. \quad (13)$$

For H_Q being Λ_b , it is zero [3]. It is a bit more complicated in the case of H_Q being $\Sigma_Q^{(*)}$. According to Ref. [7], the chromomagnetic $1/m_Q$ correction can be parametrized as

$$\bar{u}_{\Sigma_c^{(*)}}^\mu(v', s') \sigma^{\lambda\rho} \frac{1 + \not{v}'}{2} \Gamma u_{\Sigma_b^{(*)}}^\nu(v, s) M_{\lambda\rho}^{\mu\nu}, \quad (14)$$

where

$$M_{\lambda\rho}^{\mu\nu} = \frac{1}{2m_c} [\zeta_1(\omega) g_\lambda^\mu g_\rho^\nu + \zeta_2(\omega) g_\rho^\mu v'^\nu v_\lambda + \zeta_3(\omega) g_\lambda^\nu v^\mu v_\rho]. \quad (15)$$

Now consider $1/m_Q$ corrections of the current operator. The relation between the QCD currents and HQET operators is

$$\bar{c} \Gamma b = \bar{c}_{v'} \left(\Gamma - \frac{i \tilde{D}_\alpha}{2m_Q} \gamma^\alpha \Gamma \right) b_v. \quad (16)$$

For the $\Lambda_b \rightarrow \Lambda_c$ decay [15,16],

$$\bar{c}_{v'} i \tilde{D}_\alpha \Gamma b_v = \frac{\bar{\Lambda} \eta}{(1 + \omega)} \bar{u}_{\Lambda_c}(v', s') \Gamma u_{\Lambda_b}(v, s) (v_\alpha - \omega v'_\alpha), \quad (17)$$

where $\bar{\Lambda} = m_{\Lambda_Q} - m_Q$. For the $\Sigma_b^{(*)}(v, s) \rightarrow \Sigma_c^{(*)}(v', s')$ transition, the $1/m_Q$ correction is parametrized as [7]

$$\bar{c}_{v'} i \tilde{D}_\alpha \Gamma b_v = \bar{u}_{\Sigma_c^{(*)}}^\mu(v', s') \Gamma u_{\Sigma_b^{(*)}}^\nu(v, s) P_{\alpha}^{\mu\nu}, \quad (18)$$

where

$$P_{\alpha}^{\mu\nu} = \kappa_1 v'^\nu v^\mu v_\alpha + \kappa_2 v'^\nu v^\mu v'_\alpha + \kappa_3 g^{\mu\nu} v_\alpha + \kappa_4 g^{\mu\nu} v'_\alpha + \kappa_5 g_\alpha^\mu v'^\nu + \kappa_6 g_\alpha^\nu v'^\mu. \quad (19)$$

Actually, only two of these Isgur-Wise functions are independent [7],

$$\begin{aligned}\kappa_3 &= -\frac{\bar{\Sigma}}{1+\omega}\xi_1, & \kappa_4 &= \frac{\bar{\Sigma}\omega}{1+\omega}\xi_1, \\ \kappa_5 &= \bar{\Sigma}(1-\omega)\xi_2 - (\kappa_1 + \omega\kappa_2), \\ \kappa_6 &= -(\omega\kappa_1 + \kappa_2).\end{aligned}\quad (20)$$

The expressions of form factors in Eqs. (1) and (2) in terms of all these Isgur-Wise functions can be found in [7].

B. Large- N_c relations

Now consider the large- N_c limit; there are relations among the subleading order Isgur-Wise functions. The point is that the observation in Sec. I is still applicable. The only difference here is that heavy quark currents have different forms, which are irrelevant because of the heavy quark symmetry. Then in the large- N_c limit, Eq. (12) which is the charm quark kinetic energy correction should be

$$\frac{\chi(\omega)}{m_c} \left[-g_{\mu\nu} + \frac{v_\mu v'_\nu}{(1+\omega)} \right] \bar{u}_{\Sigma_c^{(*)}}^\mu(v', s') \Gamma u_{\Sigma_b^{(*)}}^\nu(v, s). \quad (21)$$

This results in the following relations:

$$\chi_1(\omega) = \chi(\omega), \quad \chi_2(\omega) = \frac{\chi(\omega)}{1+\omega}. \quad (22)$$

With the same method, we obtain a pleasant result: in the large- N_c limit, to the $1/m_Q$ order in HQET, there is no chromomagnetic corrections in the $\Sigma_b^{(*)}(v, s) \rightarrow \Sigma_c^{(*)}(v', s')$ decay,

$$\zeta_1(\omega) = \zeta_2(\omega) = \zeta_3(\omega) = 0. \quad (23)$$

This can be understood easily, since in the large- N_c limit, spins and isospins of light degrees of freedom in $\Sigma_b^{(*)}(v, s)$ and $\Sigma_c^{(*)}(v', s')$ have decoupled. This decoupling makes $\Sigma_Q^{(*)}$ no different from Λ_Q .

Therefore, in the large- N_c limit, the time-ordered product of $1/m_Q$ terms in the Lagrangian with the heavy quark current just produces a trivial correction for $\Sigma_Q^{(*)}$ decays: a redefinition of the leading order Isgur-Wise functions; this is similar to the case of Λ_Q decays.

Looking at the $1/m_c$ correction of the current operator, for $\Sigma_Q^{(*)}$ we have

$$\begin{aligned}& \bar{c}_\nu i \tilde{D}_\alpha \Gamma b_\nu \\ &= \frac{\bar{\Sigma}\eta}{(1+\omega)} (v_\alpha - \omega v'_\alpha) \\ & \quad \times \left[-g_{\mu\nu} + \frac{v_\mu v'_\nu}{1+\omega} \right] \bar{u}_{\Sigma_c^{(*)}}^\mu(v', s') \Gamma u_{\Sigma_b^{(*)}}^\nu(v, s),\end{aligned}\quad (24)$$

where $\bar{\Sigma}$ is defined as $\bar{\Sigma} = m_{\Sigma_b} - m_b \approx m_{\Sigma_c} - m_c$. Note that $\bar{\Sigma} = \bar{\Lambda}$ in the large- N_c limit. Again, we obtain some new relations as below:

$$\begin{aligned}\kappa_1 &= \frac{\bar{\Sigma}}{(1+\omega)^2}\eta, & \kappa_2 &= -\frac{\bar{\Sigma}\omega}{(1+\omega)^2}\eta, & \kappa_3 &= -\frac{\bar{\Sigma}}{1+\omega}\eta, \\ \kappa_4 &= \frac{\bar{\Sigma}\omega}{1+\omega}\eta, & \kappa_5 &= \kappa_6 = 0.\end{aligned}\quad (25)$$

It is observed that, after taking the large- N_c approximation, the relations obtained in HQET, such as Eq. (20), still hold.

C. General form factors

Up to now, we have derived all of the large- N_c relations for the $1/m_c$ corrections; $1/m_b$ corrections can be obtained similarly. Including all $1/m_Q$ corrections, in the large- N_c limit, the general form factors in Eqs. (1) and (2) are expressed as

$$\begin{aligned}F_1 &= \eta'(\omega) + \eta'(\omega) \left[\frac{\bar{\Lambda}}{2m_c} + \frac{\bar{\Lambda}}{2m_b} \right], \\ G_1 &= \eta'(\omega) - \eta'(\omega) \left[\frac{\bar{\Lambda}}{2m_c} + \frac{\bar{\Lambda}}{2m_b} \right] \left(\frac{1-\omega}{1+\omega} \right) \\ F_2 &= -\frac{\bar{\Lambda}}{m_c} \left(\frac{1}{1+\omega} \right) \eta'(\omega), \\ G_2 &= -\frac{\bar{\Lambda}}{m_c} \left(\frac{1}{1+\omega} \right) \eta'(\omega) \\ F_3 &= -\frac{\bar{\Lambda}}{m_b} \left(\frac{1}{1+\omega} \right) \eta'(\omega), & G_3 &= \frac{\bar{\Lambda}}{m_b} \left(\frac{1}{1+\omega} \right) \eta'(\omega) \\ F'_1 &= -\frac{1}{3} \eta'(\omega) - \frac{1}{3} \eta'(\omega) \left[\frac{\bar{\Sigma}}{2m_c} + \frac{\bar{\Sigma}}{2m_b} \right], \\ G'_1 &= -\frac{1}{3} \eta'(\omega) + \frac{1}{3} \eta'(\omega) \left[\frac{\bar{\Sigma}}{2m_c} + \frac{\bar{\Sigma}}{2m_b} \right] \left(\frac{1-\omega}{1+\omega} \right) \\ F'_2 &= \frac{4\eta'(\omega)}{3(1+\omega)} + \frac{\eta'(\omega)}{3(1+\omega)} \left[-\frac{\bar{\Sigma}}{m_c} + \frac{2\bar{\Sigma}}{m_b} \right], \\ G'_2 &= \frac{\bar{\Sigma}}{3m_c} \left(\frac{1}{1+\omega} \right) \eta'(\omega) \\ F'_3 &= \frac{4\eta'(\omega)}{3(1+\omega)} + \frac{\eta'(\omega)}{3(1+\omega)} \left[\frac{2\bar{\Sigma}}{m_c} - \frac{\bar{\Sigma}}{m_b} \right], \\ G'_3 &= -\frac{\bar{\Sigma}}{3m_b} \left(\frac{1}{1+\omega} \right) \eta'(\omega) \\ N'_1 &= \frac{-2\eta'(\omega)}{\sqrt{3}(1+\omega)} + \frac{-\eta'(\omega)}{\sqrt{3}(1+\omega)} \left[\frac{\bar{\Sigma}}{m_c} + \frac{\bar{\Sigma}}{m_b} \right], & K'_1 &= 0 \\ N'_2 &= 0, & K'_2 &= \frac{2}{\sqrt{3}} \eta'(\omega) \frac{\bar{\Sigma}}{m_c} \left(\frac{1}{1+\omega} \right)^2, \\ N'_3 &= \frac{2\eta'(\omega)}{\sqrt{3}(1+\omega)} + \frac{\eta'(\omega)}{\sqrt{3}(1+\omega)} \left[\frac{\bar{\Sigma}}{m_c} + \frac{\bar{\Sigma}}{m_b} \right], \\ K'_3 &= \frac{-2\eta'(\omega)}{\sqrt{3}(1+\omega)} + \frac{\eta'(\omega)}{\sqrt{3}(1+\omega)} \left[\frac{\bar{\Sigma}}{m_c} \left(\frac{1-\omega}{1+\omega} \right) - \frac{\bar{\Sigma}}{m_b} \right], \\ N'_4 &= \frac{-2\eta'(\omega)}{\sqrt{3}} - \frac{1}{\sqrt{3}} \eta'(\omega) \left[\frac{\bar{\Sigma}}{m_c} + \frac{\bar{\Sigma}}{m_b} \right], \\ K'_4 &= \frac{2\eta'(\omega)}{\sqrt{3}} - \frac{1}{\sqrt{3}} \eta'(\omega) \left[\frac{\bar{\Sigma}}{m_c} + \frac{\bar{\Sigma}}{m_b} \right] \left(\frac{1-\omega}{1+\omega} \right),\end{aligned}\quad (26)$$

where

$$\eta'(\omega) \equiv \eta(\omega) + \chi(\omega) \left(\frac{1}{m_c} + \frac{1}{m_b} \right), \quad (27)$$

and all the form factors should be multiplied by $C(\mu)$ in Eq. (3). We have checked that all the results are consistent with [7] where all $1/m_Q$ form factors are listed. After taking the large- N_c limit, all the relations of $1/m_Q$ form factors shown in [7] still hold, especially the following normalization relations at zero recoil point:

$$\begin{aligned} F_1(1) + F_2(1) + F_3(1) &= C(\mu), & G_1(1) &= C(\mu); \\ F'_1(1) + F'_2(1) + F'_3(1) &= C(\mu), & G'_1(1) &= -\frac{1}{3}C(\mu), \\ K'_4(1) &= \frac{2}{\sqrt{3}}C(\mu). \end{aligned} \quad (28)$$

In fact, through our analysis, it is easy to see that the large- N_c limit and HQET are commutative; in other words, the large- N_c approximation preserves all relations obtained in HQET.

$$\begin{aligned} \frac{d\Gamma_2(\omega)}{d\omega} &= \frac{G_F^2 |V_{cb}|^2 m_{\Omega_b}^5 r_3^3}{72\pi^3} \sqrt{(\omega^2 - 1)} \{ (\omega - 1)\kappa_3 [N'_4 - 2(\omega + 1)N'_1]^2 + (\omega + 1)\kappa_3 [K'_4 - 2(\omega - 1)K'_1]^2 \\ &+ 2(\omega + 1)[(\omega - 1)(r_3 + 1)K'_1 + (\omega^2 - 1)(K'_3 + r_3K'_2) + (\omega - r_3)K'_4]^2 + 2(\omega - 1)[(\omega + 1)(r_3 - 1)N'_1 \\ &+ (\omega^2 - 1)(N'_3 + r_3N'_2) + (\omega - r_3)N'_4]^2 + 3\kappa_3 [(\omega + 1)K'^2_4 + (\omega - 1)N'^2_4] \} \end{aligned} \quad (30)$$

where $r_3 = m_{\Omega_c}/m_{\Omega_b}$ and $\kappa_3 = 1 + r_3^2 - 2r_3\omega$.

The form factors have been expanded in Eq. (26) to the order of $1/m_c$ and $1/m_b$. There are only one Isgur-Wise function $\eta(\omega)$ at the leading order, one $\chi(\omega)$ at the subleading order, and the mass parameter $\bar{\Omega} \equiv \bar{\Sigma}$. With Eq. (8) and $\chi(\omega) \simeq 0$ [14], we obtain the decay widths as

$$\begin{aligned} \Gamma(\Omega_b \rightarrow \Omega_c l \bar{\nu}) &= 3.38 \times 10^{-14} \text{ GeV}; \\ \Gamma(\Omega_b \rightarrow \Omega_c^* l \bar{\nu}) &= 3.34 \times 10^{-14} \text{ GeV}. \end{aligned} \quad (31)$$

In the calculations, we have taken the following parameters [21]:

$$\begin{aligned} m_{\Omega_b} &= 6.07 \text{ GeV}, & m_{\Omega_c} &= 2.70 \text{ GeV}, \\ m_{\Omega_c^*} &= 2.77 \text{ GeV}, & |V_{cb}| &= 40.9 \times 10^{-3}. \end{aligned} \quad (32)$$

The pole masses of heavy quarks have been taken as $m_b = 4.83 \text{ GeV}$, $m_c = 1.43 \text{ GeV}$.

IV. DISCUSSIONS

In this paper, we have studied $O(1/m_Q)$ universal baryonic Isgur-Wise functions in the large- N_c limit; our results are explicitly listed in Eqs. (22), (23), and (25). As an application, we have calculated the semileptonic decays of Ω_b . Actually, the same results would be obtained using the

III. THE WEAK DECAYS

As an application, we now calculate $\Omega_b \rightarrow \Omega_c^{(*)}$ weak decay rates [17]. Since $\Sigma_b^{(*)}$ has the strong interaction decay mode, we mainly consider the semileptonic decays of Ω_b . In the SU(3) light quark flavor symmetry limit, $\Omega_{b(c)}^{(*)}$ baryons are identical to $\Sigma_{b(c)}^{(*)}$ baryons. Therefore, for the Isgur-Wise functions, the same results for $\Omega_{b(c)}^{(*)}$ can be obtained.

Neglecting the lepton masses, for the decay of $\Omega_b \rightarrow \Omega_c l \bar{\nu}$, the differential decay rate can be expressed [18,19] in terms of the general form factors in Eq. (1) as

$$\begin{aligned} \frac{d\Gamma_1(\omega)}{d\omega} &= \frac{G_F^2 |V_{cb}|^2 m_{\Omega_b}^5 r_2^3}{24\pi^3} \sqrt{(\omega^2 - 1)} \{ 2(\omega - 1)\kappa_2 F_1^2 \\ &+ (\omega - 1)[(1 + r_2)F'_1 + (\omega + 1)(r_2 F'_2 + F'_3)]^2 \\ &+ 2(\omega + 1)\kappa_2 G_1^2 + (\omega + 1)[(1 - r_2)G'_1 \\ &- (\omega - 1)(r_2 G'_2 + G'_3)]^2 \} \end{aligned} \quad (29)$$

where $r_2 = m_{\Omega_c}/m_{\Omega_b}$ and $\kappa_2 = 1 + r_2^2 - 2r_2\omega$.

For the decay of $\Omega_b \rightarrow \Omega_c^* l \bar{\nu}$, we have [20]

leading order large- N_c analysis in [9–11], while our method is a lot simpler.

Let us now consider the uncertainties of our results. Since the $1/m_Q$ corrections have been included, the uncertainties brought about by HQET have been suppressed to $O(\Lambda_{\text{QCD}}^2/m_c^2 \sim 1/25)$. The remaining uncertainties come from two approximations: flavor SU(3) symmetry and the large- N_c limit. Effects of flavor SU(3) violation might not be huge especially near the zero recoil point. Since Ω_b has strange quarks, we could expect the effects as [3]

$$|\xi_1(\omega) - \eta(\omega)| \sim \ln \left(\frac{m_K^2}{\mu^2} \right) \quad (33)$$

which would be small by choosing some appropriate renormalization scale μ . Then, the main uncertainties are produced by the large- N_c approximation, while we have so little knowledge about them. Sometimes they can be as large as 30%. In our case, however, it is unnecessary to be that pessimistic. As a general experience, the large- N limit is a good approximation for baryons and also good for Isgur-Wise functions. Because of replacing $\bar{\Sigma}$ with $\bar{\Omega}$ in the decay calculation, we have already taken part of the flavor SU(3) violation effects and part of the corrections to the large- N_c limit into account.

Finally, it is important to notice that, whether or not the large- N_c limit can be treated as a good approximation, at least in the vicinity of the zero recoil point, the uncertainties produced by the large- N_c limit should not be huge, since it preserves the normalizations of Isgur-Wise functions as in HQET, just like we have emphasized in Sec. III, which will be tested at the LHC or the proposed Z-factory in the near future.

ACKNOWLEDGMENTS

This work was supported in part by the National Natural Science Foundation of China under Grants No. 11075193 and No. 10821504, and by the National Basic Research Program of China under Grant No. 2010CB833000.

-
- [1] N. Isgur and M. B. Wise, *Phys. Lett. B* **232**, 113 (1989); **237**, 527 (1990); M. A. Shifman and M. B. Voloshin *Yad. Fiz.* **45**, 463 (1987) [*Sov. J. Nucl. Phys.* **45**, 292 (1987)]; M. A. Shifman and M. B. Voloshin, *Yad. Fiz.* **47**, 801 (1988) [*Sov. J. Nucl. Phys.* **47**, 511 (1988)]; E. Eichten and B. R. Hill, *Phys. Lett. B* **234**, 511 (1990); E. V. Shuryak, *Phys. Lett.* **93B**, 134 (1980).
- [2] H. Georgi, *Phys. Lett. B* **240**, 447 (1990); A. F. Falk, H. Georgi, B. Grinstein, and M. B. Wise, *Nucl. Phys.* **B343**, 1 (1990).
- [3] N. Isgur and M. B. Wise, *Nucl. Phys.* **B348**, 276 (1991); H. Georgi, *Nucl. Phys.* **B348**, 293 (1991); T. Mannel, W. Roberts, and Z. Ryzak, *Nucl. Phys.* **B355**, 38 (1991).
- [4] For reviews, see A. V. Manohar and M. B. Wise, *Heavy Quark Physics* (Cambridge University Press, Cambridge, England, 2000); A. Grozin, *Heavy Quark Effective Theory* (Springer, New York, 2004).
- [5] G. 't Hooft, *Nucl. Phys.* **B72**, 461 (1974); E. Witten, *Nucl. Phys.* **B160**, 57 (1979).
- [6] N. Isgur and M. B. Wise, *Nucl. Phys.* **B348**, 276 (1991); H. Georgi, *Nucl. Phys.* **B348**, 293 (1991); T. Mannel, W. Roberts, and Z. Ryzak, *Nucl. Phys.* **B355**, 38 (1991).
- [7] C. G. Boyd and D. E. Brahm, *Phys. Lett. B* **254**, 468 (1991).
- [8] R. F. Dashen and A. V. Manohar, *Phys. Lett. B* **315**, 425 (1993); **315**, 438 (1993); E. Jenkins, *Phys. Lett. B* **315**, 431 (1993); **315**, 441 (1993); **315**, 447 (1993); R. F. Dashen, E. Jenkins, and A. V. Manohar, *Phys. Rev. D* **49**, 4713 (1994); **51**, 3697 (1995); C. Carone, H. Georgi, and S. Osofsky, *Phys. Lett. B* **322**, 227 (1994); M. A. Luty and J. March-Russell, *Nucl. Phys.* **B426**, 71 (1994); J.-L. Gervais and B. Sakita, *Phys. Rev. D* **30**, 1795 (1984).
- [9] C. K. Chow, *Phys. Rev. D* **51**, 1224 (1995); **54**, 873 (1996).
- [10] F. Hussain and G. Thompson, *Phys. Lett. B* **335**, 205 (1994).
- [11] F. Hussain, J. G. Körner, M. Krämer, and G. Thompson, *Z. Phys. C* **51**, 321 (1991); J. G. Körner, M. Krämer, and D. Pirjol, *Prog. Part. Nucl. Phys.* **33**, 787 (1994); J. G. Körner, *Nucl. Phys. B, Proc. Suppl.* **50**, 140 (1996); F. Hussain, J. G. Körner, J. Landgraf, and S. Tawfiq, *Z. Phys. C* **69**, 655 (1996).
- [12] E. Jenkins, A. V. Manohar, and M. B. Wise, *Nucl. Phys.* **B396**, 27 (1993); C. G. Callan, Jr. and I. R. Klebanov, *Nucl. Phys.* **B262**, 365 (1985); M. Rho, D. O. Riska, and N. N. Scoccola, *Phys. Lett. B* **251**, 597 (1990); *Z. Phys. A* **341**, 343 (1992); Y. Oh, D.-P. Min, M. Rho, and N. N. Scoccola, *Nucl. Phys.* **A534**, 493 (1991).
- [13] E. Jenkins, A. V. Manohar, and M. B. Wise, *Nucl. Phys.* **B396**, 38 (1993).
- [14] J.-P. Lee, C. Liu, and H. S. Song, *Phys. Rev. D* **58**, 014013 (1998); C. Liu, *Phys. Rev. D* **57**, 1991 (1998).
- [15] M. Luke, *Phys. Lett. B* **252**, 447 (1990).
- [16] H. Georgi, B. Grinstein, and M. Wise, *Phys. Lett. B* **252**, 456 (1990).
- [17] M.-k. Du and C. Liu, *Phys. Rev. D* **84**, 056007 (2011).
- [18] Y.-B. Dai, C.-S. Huang, C. Liu, and C.-D. Lü, *Phys. Lett. B* **371**, 99 (1996).
- [19] J. G. Körner, M. Krämer, and D. Pirjol, *Prog. Part. Nucl. Phys.* **33**, 787 (1994).
- [20] A. K. Leibovich and I. W. Stewart, *Phys. Rev. D* **57**, 5620 (1998).
- [21] J. Beringer *et al.* (Particle Data Group), *Phys. Rev. D* **86**, 010001 (2012).